

# Status report of air quality in Europe for year 2024, using validated and up-to-date data



**Authors:**

Jaume Targa (4sfera Innova), María Colina (4sfera Innova), Lorena Banyuls (4sfera Innova), Alberto González Ortiz (EEA), Joana Soares (NILU)



Cover design: EEA  
Cover image: © Jaume Targa  
Layout: EEA / ETC HE

**Publication Date:** 9 April 2025

Version: 1

**DOI:** <https://doi.org/10.5281/zenodo.15168515>

#### **Legal notice**

Preparation of this report has been funded by the European Environment Agency (EEA) as part of a grant with the European Topic Centre on Human health and the environment (ETC-HE) and expresses the views of the authors. The contents of this publication do not necessarily reflect the position or opinion of the European Commission or other institutions of the European Union. Neither the European Environment Agency nor the European Topic Centre on Human health and the environment is liable for any consequence stemming from the reuse of the information contained in this publication.

#### *How to cite this report:*

Targa, J., Colina, M., Banyuls, L., González Ortiz, A., Soares, J. (2024). Status report of air quality in Europe for year 2024, using validated and up-to-date data (ETC-HE Report 2025/1). European Topic Centre on Human health and the environment.

The report is available from <https://www.eionet.europa.eu/etcs/all-etc-reports>.

**EEA activity no. 3** Human health and the environment.

**ETC-HE coordinator:** NILU - Stiftelsen NILU, Kjeller, Norway

**ETC-HE consortium partners:** Federal Environment Agency/Umweltbundesamt (UBA), Aether Limited, Czech Hydrometeorological Institute (CHMI), Institut National de l'Environnement Industriel et des Risques (INERIS), Swiss Tropical and Public Health Institute (Swiss TPH), Universitat Autònoma de Barcelona (UAB), Vlaamse Instelling voor Technologisch Onderzoek (VITO), 4sfera Innova S.L.U., klarFAKTe.U

#### **Copyright notice**

© European Topic Centre on Human health and the environment, 2025.

Reproduction is authorized provided the source is acknowledged. [Creative Commons Attribution 4.0 (International)]

More information on the European Union is available on the Internet (<http://europa.eu>).

European Topic Centre on  
Human health and the environment (ETC-HE)  
<https://www.eionet.europa.eu/etcs/etc-he>

## Acknowledgements

This report has been produced by the European Topic Centre on Human health and the Environment (ETC HE) in close cooperation with the EEA.

The ETC task manager was Jaume Targa, closely assisted by María Colina and Joana Soares, and the EEA project manager was Alberto González Ortiz closely assisted by Silvia Monge and Ragnhildur Finnbjörnsdóttir.

Additional EEA contributors were Artur Gsell. Additional ETC HE contributors were Michel Houssiau (under 4sfera Innova), Jaume Archilés (4sfera Innova) and Rune Ødegård (NILU).

Thanks are due to the air quality data suppliers in the reporting countries for collecting and providing the data on which this report is based.

## Contents

<b>1</b>	<b>Summary</b>	<b>4</b>
1.1	Particulate matter . . . . .	6
1.2	Ozone . . . . .	6
1.3	Nitrogen dioxide . . . . .	6
1.4	Sulphur dioxide . . . . .	6
1.5	Editorial note . . . . .	8
<b>2</b>	<b>Introduction</b>	<b>9</b>
<b>3</b>	<b>Status of particulate matter ambient air concentrations</b>	<b>14</b>
3.1	Status of PM <sub>10</sub> concentrations . . . . .	14
3.2	Status of PM <sub>2.5</sub> concentrations . . . . .	24
<b>4</b>	<b>Status of ozone ambient air concentrations</b>	<b>31</b>
<b>5</b>	<b>Status of nitrogen dioxide ambient air concentrations</b>	<b>41</b>
<b>6</b>	<b>Status of sulphur dioxide ambient air concentrations</b>	<b>48</b>
<b>7</b>	<b>Abbreviations, units and symbols</b>	<b>52</b>
<b>8</b>	<b>Annex 1</b>	<b>53</b>
<b>9</b>	<b>Annex 2 - Air quality status in relation to the new standards in the revised AAQD (EU)</b>	
	<b>2024/2081</b>	<b>57</b>
9.1	Status of PM <sub>10</sub> concentrations . . . . .	60
9.2	Status of PM <sub>2.5</sub> concentrations . . . . .	63
9.3	Status of O <sub>3</sub> ambient air concentrations . . . . .	66
9.4	Status of NO <sub>2</sub> ambient air concentrations . . . . .	68
9.5	Status of SO <sub>2</sub> concentrations . . . . .	71



## 1 Summary

The *2024 Status report of air quality in Europe* presents summarized information on the air quality data for the protection of health reported in the previous years. The reported 2024 monitoring data used in this analysis was reported as up-to-date (UTD) data, prior to final quality control and validated data reporting by the countries <sup>(1)</sup>. It provides information on the following pollutants, regulated by the Ambient Air Quality Directives (AAQD) (EU, 2008, 2024)

- PM<sub>10</sub>: Particulate matter with a diameter of 10 µm or less
- PM<sub>2.5</sub>: Particulate matter with a diameter of 2.5 µm or less
- O<sub>3</sub>: Tropospheric ozone
- NO<sub>2</sub>: Nitrogen dioxide
- SO<sub>2</sub>: Sulphur dioxide

It also offers a comparison with the situation in previous years. For those years, validated data are considered.

Data included in this report was received by 19 February 2025 from the reporting countries. By that date the reporting status of 2024 up-to-date data is summarized in Figure 1, where a green box indicates that the referred pollutant was reported by the referred country and a grey box indicates the contrary (that the referred pollutant was not reported by the referred country). Please see editorial notes at the end of this Chapter on additional information on the data used. The number of stations by country reporting each pollutant, with the minimum data coverage for at least one of the aggregations used in the report, is also included in Figure 1, while Table 3 in the Annex summarizes the number of stations, with the minimum data coverage for at least one of the aggregations used in the report, at different country aggregations. Data from stations that do not fulfil the criteria from Box 1.1 are excluded from this report. Please be aware that the number of stations presented in Figure 1 and Table 3, that corresponds to all reported stations fulfilling the minimum data coverage criteria for at least one of the aggregations used in the report, may be different to the one presented in the corresponding boxplots, as there could be some stations not fulfilling the minimum data coverage criteria for the corresponding aggregation.

---

<sup>1</sup><https://eeadmz1-cws-wp-air02-dev.azurewebsites.net/aq-ereporting/>

Figure 1: Number of stations, for each country and each pollutant, that in 2024 reported data with the minimum data coverage for at least one of the aggregations used in the report, by 19 February 2025

	PM10	PM2.5	O3	NO2	SO2
Albania -					
Andorra -	1	1	2	1	1
Austria -	116	54	106	138	57
Belgium -	69	71	33	77	19
Bosnia and Herzegovina -					
Bulgaria -	22	3	20	21	24
Croatia -	12	10	14	14	9
Cyprus -	2	1	3	3	3
Czechia -	75	48	55	59	41
Denmark -	2		8	10	1
Estonia -	6	7	9	9	9
Finland -	32	19	15	31	13
France -	328	230	291	360	77
Germany -	369	322	277	502	94
Greece -	20	10	19	18	8
Hungary -	22	10	16	21	18
Iceland -	7	5		8	11
Ireland -	19	16	9	16	7
Italy -	431	224	292	530	153
Kosovo -	12	12	10	9	10
Latvia -	8	5	7	7	4
Liechtenstein -					
Lithuania -	14	6	13	14	10
Luxembourg -	3	3	5	9	1
Malta -	5	5	4	5	4
Montenegro -					
Netherlands -	65	24	40	70	13
North Macedonia -	12	11	17	19	14
Norway -	50	47	12	39	5
Poland -	175	89	99	129	83
Portugal -	58	20	46	62	18
Romania -	44	12	23	20	29
Serbia -		1	8	1	
Slovakia -	48	47	24	39	18
Slovenia -	18	18	11	11	3
Spain -	390	238	433	520	398
Sweden -	49	27	22	40	
Switzerland -	30	9	30	31	8
Türkiye -					

The countries included in this report and that, therefore, appear in Figure 1, are those with the obligation to report data under the AAQD or that have voluntary reported data. These countries are the EU-27 (Austria, Belgium, Bulgaria, Croatia, Cyprus, Czechia, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, the Netherlands, Poland, Portugal, Romania, Slovakia, Slovenia, Spain and Sweden); the five other member countries of the EEA (Iceland, Liechtenstein, Norway, Switzerland and Türkiye) that, together with the EU-27 form the EEA-32; the six EEA's cooperating countries from the Western Balkans (Albania, Bosnia and Herzegovina, Kosovo under UN Security Council Resolution 1244/99, Montenegro, North Macedonia and Serbia) that, together with the EEA-32 form the EEA-38; and the voluntary reporting country of Andorra.

The air quality data are stored at the EEA's e-reporting database <sup>(2)</sup>. Therefore, this is the source for all maps and figures in the report. UTD data is stored temporarily until it is replaced by CDR data.

### 1.1 Particulate matter

For PM<sub>10</sub>, concentrations above the EU daily limit value (50 µg/m<sup>3</sup>) were registered at 7 % of the reporting stations. These stations were in 13 countries in EU-27 and in 2 other reporting countries. For PM<sub>2.5</sub>, concentrations above the EU annual limit value (25 µg/m<sup>3</sup>) were registered at 0.6 % of the reporting stations. These stations were in 2 countries in EU-27 and in 1 other reporting countries.

The long-term World Health Organization air quality guideline (WHO AQG) level for PM<sub>10</sub> (15 µg/m<sup>3</sup>) was exceeded at 59 % of the stations in 27 countries of the EU-27 and 4 other reporting countries. The long-term WHO AQG level for PM<sub>2.5</sub> (5 µg/m<sup>3</sup>) was exceeded at 93 % of the stations located in 25 countries of the EU-27 and 6 other reporting countries.

### 1.2 Ozone

13 % of stations registered concentrations above the EU target value for O<sub>3</sub> (120 µg/m<sup>3</sup>) for the protection of human health. These stations were located in 16 countries of the EU-27 and 3 other reporting countries. The long-term EU objective (120 µg/m<sup>3</sup>) was met in only 22 % of the stations. The short-term WHO AQG level for O<sub>3</sub> (100 µg/m<sup>3</sup>) was exceeded in 92 % of all the reporting stations, and concentrations above the long-term WHO AQG level for O<sub>3</sub> (60 µg/m<sup>3</sup>) were registered in 98 % of all reporting stations.

### 1.3 Nitrogen dioxide

Around 1 % of all the reporting stations recorded concentrations above the EU annual limit value for NO<sub>2</sub> (40 µg/m<sup>3</sup>). These stations were located in 6 countries of the EU-27 and 1 other reporting country. 96 % of concentrations above this limit value were observed at traffic stations.

On the contrary, 65 % of stations, located in 27 countries of the EU-27 and 7 other reporting countries reported concentrations above the WHO AQG level of 10 µg/m<sup>3</sup>.

### 1.4 Sulphur dioxide

For SO<sub>2</sub>, regarding the EU daily limit value (125 µg/m<sup>3</sup>), concentrations above were registered at 0.4 % of the reporting stations. These stations were in 2 country of the EU-27 and 2 other reporting countries . On the other hand, concerning the daily WHO AQG level (40 µg/m<sup>3</sup>), 3 %

---

<sup>2</sup>[https://discomap.eea.europa.eu/App/AQViewer/index.html?fqn=Airquality\\_Dissem.b2g.AirQualityStatistics](https://discomap.eea.europa.eu/App/AQViewer/index.html?fqn=Airquality_Dissem.b2g.AirQualityStatistics)

of all reporting SO<sub>2</sub> stations measured SO<sub>2</sub> concentrations above. These stations were located in 10 countries of the EU-27 and 2 other reporting countries .

## 1.5 Editorial note

Values in Table 4 in Annex 1 are considered outliers and were not taken into account for the analysis presented in this report.

On 20 November 2024, the revised Directive (EU) 2024/2881 (EU, 2024) on ambient air quality and cleaner air for Europe was published and it entered into force on 10 December 2024. It sets new or revised air quality standards to be reached by 1 January 2030. The main analysis in this report is done against the AQ standards defined in the 2004 and 2008 AAQD (EU, 2008, 2024), which are the ones currently applicable. Annex 2 benchmarks the situation in year 2024 with respect to some of the new and/or revised AQ standards, as an analysis of the 'distance to target' from the 2024 status.

Once the production of the report was closed, North Macedonia informed the EEA that some SO<sub>2</sub> data from station MK0044A Kavadarci were wrongly reported. As a result of removing those wrong data, concentrations in that station would not be above the SO<sub>2</sub> daily limit value.

## 2 Introduction

The *2024 Status report of air quality in Europe* presents summarized information on the air quality data reported up to 2024. The 2024 data was reported as up-to-date (UTD) data in a continuous basis prior to final quality control and official reporting of validated data by the countries, which will be done under the 2025 September reporting cycle (validated assessment data for 2024, deadline of submission 30 September 2025). This report aims at informing on the current status of ambient air quality in Europe, based on the most updated data available for the analysis of a complete calendar year. Furthermore, it informs on progress towards meeting the air quality standards established for the protection of health in the Ambient Air Quality Directives (EU, 2004, 2008) (Table 1) and the World Health Organization air quality guideline levels (WHO, 2000, 2006, 2021) (Table 2)<sup>(3)</sup>.

This report builds on the former EEA “Air quality in Europe report” (EEA, 2020) content, figures and maps regarding the status of monitored air quality in Europe. The report focuses on the analysis of the main pollutants, to allow a meaningful preliminary analysis of their concentration status in Europe. It provides:

- a European overview of the monitoring stations that reported UTD 2024 data, and of their concentrations in relation to the EU legal standards set in the 2008 EU Ambient Air Quality Directive (EU, 2008) and WHO AQG levels for each pollutant;
- a map with the 2024 UTD concentrations at station level for each pollutant;
- a boxplot graph summarizing for each country the range of concentrations (highlighting the lowest, highest, average and the 25 and 75 percentiles) for PM<sub>10</sub>, PM<sub>2.5</sub>, NO<sub>2</sub> and O<sub>3</sub>.

Furthermore, it provides:

- maps with the situation at station level for the previous three years (using validated data). In this way, any significant change in the spatial distribution of the values above the set thresholds in the legends can be observed (assuming the UTD stations dataset is complete);

---

<sup>3</sup>Nevertheless, in this report the following standards and guideline levels are not analysed: exposure concentration obligation and national exposure reduction target for PM<sub>2.5</sub>, information and alert thresholds for O<sub>3</sub>, alert threshold for NO<sub>2</sub>, annual target value for BaP, alert threshold for SO<sub>2</sub>, limit value for CO maximum daily 8-hour mean, annual limit value for C<sub>6</sub>H<sub>6</sub>, annual limit value for Pb, target value for As, target value for Cd, and target value for Ni in Table 1; and hourly air quality guideline level for NO<sub>2</sub>, reference level for annual mean of BaP, 10 minutes air quality guideline level for SO<sub>2</sub>, air quality guideline level for CO, reference level for annual mean of C<sub>6</sub>H<sub>6</sub>, air quality guideline level for Pb, reference level for annual mean of As, air quality guideline level for Cd, and reference level for annual mean of Ni in Table 2.

- heatmaps with the evolution of the mean and the maximum measured concentrations at country level since 2000 (or since when available, using validated data for all years up to 2023).

Please be aware that the number of stations can vary once the validated dataset for 2024 is received by 30 September 2025. In some figures like the boxplots, the final order of the countries may vary once the validated data are submitted.

**Table 1: Air quality standards for the protection of health, as given in the EU 2004 and 2008 Ambient Air Quality Directives**

Pollutant	Averaging period	Legal nature and concentration	Comments
PM <sub>10</sub>	1 day	Limit value: 50 µg/m <sup>3</sup>	Not to be exceeded on more than 35 days per year
	Calendar year	Limit value: 40 µg/m <sup>3</sup>	
PM <sub>2,5</sub>	Calendar year	Limit value: 25 µg/m <sup>3</sup>	Stage 1
		Indicative limit value: 20 µg/m <sup>3</sup>	Stage 2: indicative limit value to be reviewed by the Commission in 2013. It remained unchanged after that revision
		Exposure concentration obligation: 20 µg/m <sup>3</sup>	Average Exposure Indicator (AEI) <sup>(a)</sup> in 2015 (2013-2015 average)
		National Exposure reduction target: 0-20 percentage reduction in exposure	AEI <sup>(a)</sup> in 2020, the percentage reduction depends on the initial AEI
O <sub>3</sub>	Maximum daily 8-hour mean	Target value: 120 µg/m <sup>3</sup>	Not to be exceeded on more than 25 days/year, averaged over 3 years <sup>(b)</sup>
		Long term objective: 120 µg/m <sup>3</sup>	
	1 hour	Information threshold: 180 µg/m <sup>3</sup> Alert threshold: 240 µg/m <sup>3</sup>	
NO <sub>2</sub>	1 hour	Limit value: 200 µg/m <sup>3</sup>	Not to be exceeded on more than 18 hours per year
		Alert threshold: 400 µg/m <sup>3</sup>	To be measured over 3 consecutive hours over 100 km <sup>2</sup> or an entire zone
	Calendar year	Limit value: 40 µg/m <sup>3</sup>	
BaP	Calendar year	Target value: 1 ng/m <sup>3</sup>	Measured as content in PM <sub>10</sub>
SO <sub>2</sub>	1 hour	Limit value: 350 µg/m <sup>3</sup>	Not to be exceeded on more than 24 hours per year
		Alert threshold: 500 µg/m <sup>3</sup>	To be measured over 3 consecutive hours over 100 km <sup>2</sup> or an entire zone
	1 day	Limit value: 125 µg/m <sup>3</sup>	Not to be exceeded on more than 3 days per year
CO	Maximum daily 8-hour mean	Limit value: 10 mg/m <sup>3</sup>	
C <sub>6</sub> H <sub>6</sub>	Calendar year	Limit value: 5 µg/m <sup>3</sup>	
Pb	Calendar year	Limit value: 0.5 µg/m <sup>3</sup>	Measured as content in PM <sub>10</sub>
As	Calendar year	Target value: 6 ng/m <sup>3</sup>	Measured as content in PM <sub>10</sub>
Cd	Calendar year	Target value: 5 ng/m <sup>3</sup>	Measured as content in PM <sub>10</sub>
Ni	Calendar year	Target value: 20 ng/m <sup>3</sup>	Measured as content in PM <sub>10</sub>

**Notes:**

<sup>a</sup> AEI: based upon measurements in urban background locations established for this purpose by the Member States, assessed as a 3-year running annual mean.

<sup>b</sup> In the context of this report, only the maximum daily 8-hour means in 2024 are considered, so no average over the period 2022 - 2024 is presented.

**Sources:**

EU (2004, 2008).



**Table 2: WHO air quality guideline (AQG) levels and estimated reference levels (RL) <sup>(a)</sup>**

Pollutant	Averaging period	AQG	RL	Comments
PM <sub>10</sub>	1 day	45 µg/m <sup>3</sup>		99th percentile (3-4 exceedance days per year). Updated 2021 guideline
	Calendar year	15 µg/m <sup>3</sup>		Updated 2021 guideline
PM <sub>2.5</sub>	1 day	15 µg/m <sup>3</sup>		99th percentile (3-4 exceedance days per year). Updated 2021 guideline
	Calendar year	5 µg/m <sup>3</sup>		Updated 2021 guideline
O <sub>3</sub>	Maximum daily 8-hour mean	100 µg/m <sup>3</sup>		99th percentile (3-4 exceedance days per year). Updated 2021 guideline
	Peak season <sup>(b)</sup>	60 µg/m <sup>3</sup>		New 2021 guideline
NO <sub>2</sub>	1 hour	200 µg/m <sup>3</sup>		
	1 day	25 µg/m <sup>3</sup>		99th percentile (3-4 exceedance days per year). New 2021 guideline
	Calendar year	10 µg/m <sup>3</sup>		Updated 2021 guideline
BaP	Calendar year		0.12 ng/m <sup>3</sup>	
SO <sub>2</sub>	10 minutes	500 µg/m <sup>3</sup>		
	1 day	40 µg/m <sup>3</sup>		99th percentile (3-4 exceedance days per year). Updated 2021 guideline
CO	1 hour	30 mg/m <sup>3</sup>		
	Maximum daily 8-hour mean	10 mg/m <sup>3</sup>		
	1 day	4 mg/m <sup>3</sup>		99th percentile (3-4 exceedance days per year). New 2021 guideline
C <sub>6</sub> H <sub>6</sub>	Calendar year		1.7 µg/m <sup>3</sup>	
Pb	Calendar year	0.5 µg/m <sup>3</sup>		
As	Calendar year		6.6 ng/m <sup>3</sup>	
Cd	Calendar year	5 ng/m <sup>3</sup> <sup>(c)</sup>		
Ni	Calendar year		25 ng/m <sup>3</sup>	

**Notes:**

<sup>a</sup> As WHO has not set an AQG level for BaP, C<sub>6</sub>H<sub>6</sub>, As and Ni, the RL was estimated assuming an acceptable risk of additional lifetime cancer risk of approximately 1 in 100 000.

<sup>b</sup> Average of daily maximum 8-hour mean concentration in the six consecutive months with the highest six-month running average O<sub>3</sub> concentration.

<sup>c</sup> AQG set to prevent any further increase of Cd in agricultural soil, likely to increase the dietary intake of future generations.

**Sources:**

WHO (2000, 2006, 2021).

**Box 1.1 Classification of monitoring stations and criteria used for the assessment**

Fixed sampling points in Europe are situated at different types of stations following rules for macro- and micro-scale siting. Briefly, depending on the predominant emission sources, stations are classified as follows:

- traffic stations: located in close proximity to a single major road;
- industrial stations: located in close proximity to an industrial area or an industrial source;
- background stations: where pollution levels are representative of the average exposure of the general population or vegetation.

Depending on the distribution/density of buildings, the area surrounding the station is classified as follows:

- urban: continuously built-up urban area;
- suburban: largely built-up urban area;
- rural: all other areas.

For the pollutants considered in this report, monitoring stations have to fulfil the criterion of reporting more than 75 % of valid data out of all the possible data in a year to be included in this assessment. Reporting stations not fulfilling the minimum data coverage could be found at the [Annual AQ statistics table](#).

Measurement data are rounded following the general recommendations under (EU, 2011). The number of considered decimals are indicated in the legend of the corresponding maps.

The assessments, in the cases of PM and SO<sub>2</sub>, do not account for the fact that the Ambient Air Quality Directive (EU, 2008) provides Member States with the possibility of subtracting contributions to the measured concentrations from natural sources and winter road sanding/salting under specific circumstances.

### 3 Status of particulate matter ambient air concentrations


#### 3.1 Status of PM<sub>10</sub> concentrations

The EEA received PM<sub>10</sub> data for 2024, with sufficient valid measurements from 2508 stations for the calculation of annual mean concentrations and from 2503 stations in relation to the daily limit value. The stations were located in all the reporting countries shown in Figure 1.

Thirteen countries in EU-27, and two other reporting countries reported PM<sub>10</sub> concentrations above the EU daily limit value of 50 µg/m<sup>3</sup> (Figure 2). This was the case for 7 % (174) of reporting stations. In total, 97 % of those stations were either urban (77 %) or suburban (20 %). The stricter value of the WHO AQG level for PM<sub>10</sub> daily mean (45 µg/m<sup>3</sup>) was exceeded at 59 % (1468) of the stations in all the reporting countries, except in Denmark, Luxembourg and Switzerland (Figure 8).

Concentrations above the PM<sub>10</sub> annual limit value (40 µg/m<sup>3</sup>) were monitored in 0.6 % (14 stations) of all the reporting stations, located in 2 countries in EU-27, and 2 other reporting countries. The stricter value of the WHO AQG level for PM<sub>10</sub> annual mean (15 µg/m<sup>3</sup>) was exceeded at 59 % (1472) of the stations in all the reporting countries, except in Andorra and Switzerland (Figure 5).

Figure 2: UTD Map and boxplot of PM<sub>10</sub> concentrations in 2024 - daily limit value




Note: The graph is based, for each country, on the 90.4 percentile of daily mean concentration values corresponding to the 36th highest daily mean in a complete time series. For each country, the number of stations considered for 2024 (in brackets) are given. The boxplot represents the lowest (bottom of the whisker), highest (top of the whisker) and average (black dot) 90.4 percentile values (in µg/m³). The rectangles mark the 25th and 75th percentiles. At 25 % of the stations, levels are below the 25th percentile; at 25 % of the stations, concentrations are above the 75th percentile. The daily limit value set by EU legislation is marked by the horizontal line. The graph should be read in relation to the above map, as a country's situation depends on the number of stations considered.

Figure 3 shows the maps of the 90.4 percentile of PM<sub>10</sub> daily mean concentrations (equivalent to the PM<sub>10</sub> daily limit value) for four years. In this way, any significant change in the spatial


distribution of the values above the set thresholds in the legends can be observed. Note that only the last year's map (2024) is based on UTD data, while the previous three years are based on officially reported validated data (CDR).

*Figure 3: Maps of PM<sub>10</sub> concentrations (daily limit value) for the last 4 years*




Heatmaps with the evolution from 2000 of the mean (top) and the maximum (bottom) 90.4 percentile of PM<sub>10</sub> daily mean concentrations at country level are shown in figure 4. In this way, the evolution along years of the average and maximum measured concentration levels can be seen for each country. Note that meteorological variability has a considerable impact on year-to-year changes in ambient air concentrations of air pollutants (EEA, 2020), and the last year (2024) is based on UTD data, while the previous years are based on officially reported validated data.

Figure 4: Evolution of mean (top) and maximum (bottom) 90.4 percentile of PM<sub>10</sub> daily mean concentrations (daily limit value) per country from 2000




Note: It is important to note that the figure is not based on a consistent set of stations. The number, location and classification of the stations included may vary from year to year.

Figure 5: UTD Map and boxplot of PM<sub>10</sub> concentrations in 2024 - annual limit value



Note: Observed concentrations of PM<sub>10</sub> in 2024. The possibility of subtracting contributions to the measured concentrations from natural sources and winter road sanding/salting has not been considered. The last two colour categories indicate stations reporting concentrations above the EU annual limit value (40 µg/m<sup>3</sup>). The first colour category indicate stations reporting values below the WHO AQG level for PM<sub>10</sub> (15 µg/m<sup>3</sup>). Only stations with more than 75 % of valid data have been included in the map.




Note: The graph is based on annual mean concentration values. For each country, the number of stations considered for 2024 (in brackets) are given. The boxplot represents the lowest (bottom of the whisker), highest (top of the whisker) and average (black dot) annual mean values (in µg/m<sup>3</sup>). The rectangles mark the 25th and 75th percentiles. At 25 % of the stations, levels are below the 25th percentile; at 25 % of the stations, concentrations are above the 75th percentile. The annual limit value set by EU legislation is marked by the upper continuous horizontal line. The WHO AQG level is marked by the lower dashed horizontal line. The graph should be read in relation to the above map, as a country's situation depends on the number of stations considered.



The highest value in the boxplot, Italy ( $217 \mu\text{g}/\text{m}^3$ ), has not been included in the graph for representation purposes.

Figure 6 shows the maps of  $\text{PM}_{10}$  annual mean concentrations at station level for the last four years. In this way, any significant change in the spatial distribution of the values above the set thresholds in the legends can be observed. Note that only the last year's map (2024) is based on UTD data, while the previous three years are based on officially reported validated data (CDR).

*Figure 6: Maps of  $\text{PM}_{10}$  concentrations (annual limit value) for the last 4 years*




Maps for years before 2023 are different to the ones published in previous reports because the bands in the legend have been modified to accommodate the 2030 EU annual limit value (EU, 2024).

Heatmaps with the evolution from 2000 of the mean (top) and the maximum (bottom) annual mean  $\text{PM}_{10}$  concentrations at country level are shown in figure 7. In this way, the evolution along years of the average and maximum measured concentration levels can be seen for each country. Note that meteorological variability has a considerable impact on year-to-year changes




in ambient air concentrations of air pollutants (EEA, 2020), and the last year (2024) is based on UTD data, while the previous years are based on officially reported validated data.

*Figure 7: Evolution of mean (top) and maximum (bottom) PM<sub>10</sub> annual mean concentrations (annual limit value) per country from 2000*



Note: It is important to note that the figure is not based on a consistent set of stations. The number, location and classification of the stations included may vary from year to year.


Figure 8: UTD Map of PM<sub>10</sub> concentrations in 2024 - daily WHO AQG level



Note: Observed concentrations of PM<sub>10</sub> in 2024. The map shows the 99 percentile of the PM<sub>10</sub> daily mean concentrations, equivalent to 3–4 exceedance days per year, according to the definition of the daily WHO AQG level (45 µg/m<sup>3</sup>). The first colour category indicates stations with concentrations below this AQG level. Only stations with more than 75 % of valid data have been included in the map.


Figure 9 shows the maps of the 99 percentile of PM<sub>10</sub> daily mean concentrations (equivalent to the WHO AQG level for PM<sub>10</sub> daily mean level) for the last four years. In this way, any significant change in the spatial distribution of the values above the set thresholds in the legends can be observed. Note that only the last year's map (2024) is based on UTD data, while the previous three years are based on officially reported validated data (CDR).

*Figure 9: Maps of PM<sub>10</sub> concentrations (daily WHO AQG level) for the last 4 years*



Heatmaps with the evolution from 2013 of the mean (top) and the maximum (bottom) 99 percentile of PM<sub>10</sub> daily mean concentrations at country level are shown in figure 10. In this way, the evolution along years of the average and maximum measured concentration levels can be seen for each country. Note that meteorological variability has a considerable impact on year-to-year changes in ambient air concentrations of air pollutants (EEA, 2020), and the last year (2024) is based on UTD data, while the previous years are based on officially reported validated data.

Figure 10: Evolution of mean (top) and maximum (bottom) 99 percentile of PM<sub>10</sub> daily mean concentrations (daily WHO AQG level) per country from 2013



Note: It is important to note that the figure is not based on a consistent set of stations. The number, location and classification of the stations included may vary from year to year.

### 3.2 Status of PM<sub>2.5</sub> concentrations


Regarding PM<sub>2.5</sub>, data with sufficient valid measurements were received from 1605 stations for the calculation of annual mean concentrations and from 1597 stations in relation to the short-term WHO AQG level. These stations were located in all the reporting countries shown in Figure 1.

The PM<sub>2.5</sub> concentrations were higher than the EU annual limit value (25 µg/m<sup>3</sup>) in two countries in EU-27 and one other reporting countries (Figure 11). These concentrations above the limit value were registered in 0.6 % of all the reporting stations and occurred primarily (100 % of cases) in urban (67 %) or suburban (33 %) areas.


The WHO AQG level for PM<sub>2.5</sub> annual mean (5 µg/m<sup>3</sup>) was exceeded at 93 % of the stations, located in 31 of the 33 countries reporting PM<sub>2.5</sub> data (Figure 11). Estonia and Iceland did not report any concentrations above the WHO AQG level for PM<sub>2.5</sub>.

The WHO AQG level for PM<sub>2.5</sub> daily mean (15 µg/m<sup>3</sup>), expressed as percentile 99, it was exceeded at 96 % (1534 stations) of the stations in all the reporting countries (Figure 14).

Figure 11: UTD Map and boxplot of PM<sub>2.5</sub> concentrations in 2024 - annual limit value




Note: Observed concentrations of PM<sub>2.5</sub> in 2024. The possibility of subtracting contributions to the measured concentrations from natural sources has not been considered. The last two colour categories indicate stations reporting concentrations above the EU indicative annual limit value (20 µg/m<sup>3</sup>) or the EU annual limit value (25 µg/m<sup>3</sup>). The first colour category indicates stations reporting values below the WHO AQG level for PM<sub>2.5</sub> (5 µg/m<sup>3</sup>). Only stations with more than 75 % of valid data have been included in the map.



Note: The graph is based on annual mean concentration values. For each country, the number of stations considered for 2024 (in brackets) are given. The boxplot represents the lowest (bottom of the whisker), highest (top of the whisker) and average (black dot) annual mean values (in µg/m<sup>3</sup>). The rectangles mark the 25th and 75th percentiles. At 25 % of the stations, levels are below the 25th percentile; at 25 % of the stations, concentrations are above the 75th percentile. The annual limit value and the indicative annual limit value set by EU legislation are marked by the upper continuous horizontal lines at 25 and 20, respectively. The WHO AQG level is marked by the lower dashed horizontal line. The graph should be read in relation to the above map, as a country's situation depends on the number of stations considered.

Figure 12 shows the maps of measured PM<sub>2.5</sub> annual mean concentrations for the last four years. In this way, any significant change in the spatial distribution of the values above the set thresholds in the legends can be observed. Note that only the last year's map (2024) is based on UTD data, while the previous three years are based on officially reported validated data (CDR).


*Figure 12: Maps of PM<sub>2.5</sub> concentrations (annual limit value) for the last 4 years*



Heatmaps with the evolution from 2000 of the mean (top) and the maximum (bottom) PM<sub>2.5</sub> annual mean concentrations at country level are shown in figure 13. In this way, the evolution along years of the average and maximum measured concentration levels can be seen for each country. Note that meteorological variability has a considerable impact on year-to-year changes in ambient air concentrations of air pollutants (EEA, 2020), and the last year (2024) is based on UTD data, while the previous years are based on officially reported validated data.




Figure 13: Evolution of mean (top) and maximum (bottom) PM<sub>2.5</sub> annual mean concentrations (annual limit value) per country from 2000



Note: It is important to note that the figure is not based on a consistent set of stations. The number, location and classification of the stations included may vary from year to year.




Figure 14: UTD Map of  $PM_{2.5}$  concentrations in 2024 - daily WHO AQG level



Note: Observed concentrations of  $PM_{2.5}$  in 2024. The map shows the 99 percentile of the  $PM_{2.5}$  daily mean concentrations, equivalent to 3–4 exceedance days per year, according to the definition of the daily WHO AQG level ( $15 \mu g/m^3$ ). The first colour category indicates stations with concentrations below this AQG level. Only stations with more than 75 % of valid data have been included in the map.


Figure 15 shows the maps of the 99 percentile of PM<sub>2.5</sub> daily mean concentrations (equivalent to the WHO AQG level for PM<sub>2.5</sub> daily mean level) for the last four years. In this way, any significant change in the spatial distribution of the values above the set thresholds in the legends can be observed. Note that only the last year's map (2024) is based on UTD data, while the previous three years are based on officially reported validated data (CDR).

*Figure 15: Maps of PM<sub>2.5</sub> concentrations (daily WHO AQG level) for the last 4 years*



Heatmaps with the evolution from 2013 of the mean (top) and the maximum (bottom) 99 percentile of PM<sub>2.5</sub> daily mean concentrations at country level are shown in figure 16. In this way, the evolution along years of the average and maximum measured concentration levels can be seen for each country. Note that meteorological variability has a considerable impact on year-to-year changes in ambient air concentrations of air pollutants (EEA, 2020), and the last year (2024) is based on UTD data, while the previous years are based on officially reported validated data.

Figure 16: Evolution of mean (top) and maximum (bottom) 99 percentile of PM<sub>2.5</sub> daily mean concentrations (daily WHO AQG level) per country from 2013



Note: It is important to note that the figure is not based on a consistent set of stations. The number, location and classification of the stations included may vary from year to year.

## 4 Status of ozone ambient air concentrations

Data for O<sub>3</sub> were reported from 1938 stations for the calculation of EU standards, from 1938 stations in relation to the short-term WHO AQG level, and from 1876 stations for the long-term WHO AQG level. These stations were located in all the reporting countries shown in Figure 1.


16 countries in EU-27 and 3 other reporting countries registered concentrations above the O<sub>3</sub> target value threshold (120 µg/m<sup>3</sup>) more than 25 times this year (Figure 17). In total, 13 % of all stations reporting O<sub>3</sub> showed concentrations above the target value threshold for the protection of human health. In addition, only 22 % (420) of all stations fulfilled the long-term objective (120 µg/m<sup>3</sup>). 87 % of the stations with values above the long-term objective were background stations.

8 % (146) of all stations and only 19 of the 502 reported rural background stations had values below the short-term WHO AQG value for O<sub>3</sub> (100 µg/m<sup>3</sup>) (Figure 20), set for the protection of human health. The long-term, peak season<sup>(4)</sup>, WHO AQG level (60 µg/m<sup>3</sup>) was exceeded in 98 % (1837) of all stations located in 27 countries in EU-27 and 6 other reporting countries. Only 1 of the 484 reported rural background stations had values below this AQG level (Figure 23).


---

<sup>4</sup>The peak season is calculated for each station as the average of daily maximum 8-hour mean O<sub>3</sub> concentration in the six consecutive months with the highest six-month running-average O<sub>3</sub> concentration. That means that, for each station, twelve 6-months running averages of the daily 8-h max are calculated (1 August YY-1 to 31 January YY, ..., 1 January YY to 30 June YY, ..., 1 July YY to 31 December YY) and the maximum of those 12 values is selected as the peak season concentration. Please check also Data Dictionary - Vocabulary (<https://dd.eionet.europa.eu/vocabularyconcept/aq/aggregationprocess/P1Y-maxP6M-P8H-dmax/view?vocabularyFolider.workingCopy=false&facet=HTML+Representation>).

Figure 17: UTD Map and boxplot of O<sub>3</sub> concentrations in 2024



Note: Observed concentrations of O<sub>3</sub> in 2024. The map shows the 93.2 percentile of the O<sub>3</sub> maximum daily 8–hour mean, representing the 26th highest value in a complete series. It is related to the O<sub>3</sub> target value. At sites marked with the last two colour categories, the 26th highest daily O<sub>3</sub> concentrations were above the 120 µg/m<sup>3</sup> threshold, implying values above the target value threshold. Please note that the legal definition of the target value considers not only 1 year but the average over 3 years. Only stations with more than 75 % of valid data have been included in the map.




Note: The graph is based, for each country, on the 93.2 percentile of the maximum daily 8–hour mean concentration values, corresponding to the 26th highest daily maximum of the running 8–hour mean in a complete time series. For each country, the number of stations considered for 2024 (in brackets) are given. The boxplot represents the lowest (bottom of the whisker), highest (top of the whisker) and average (black dot) values (in µg/m<sup>3</sup>). The rectangles mark the 25th and 75th percentiles. At 25 % of the stations, levels are below the 25th percentile; at 25 % of the stations, concentrations are above the 75th percentile. The target value threshold set by the EU legislation is marked by the horizontal line. Please note that the legal definition of the target value considers not only 1 year but the average over 3 years. The graph should be read in relation to the above map, as a country's situation depends on the number of stations considered.

The highest value in the boxplot, Italy (166 µg/m<sup>3</sup>), has not been included in the graph for representation purposes.

Figure 18 shows the maps of the observed 93.2 percentile of the O<sub>3</sub> maximum daily 8-hour mean concentrations (O<sub>3</sub> target value) for the last four years. In this way, any significant change in the spatial distribution of the values above the set thresholds in the legends can be observed. Note that only the last year's map (2024) is based on UTD data, while the previous three years are based on officially reported validated data (CDR).


*Figure 18: Maps of O<sub>3</sub> concentrations (related to the target value) for the last 4 years*



Note: Please be aware that the TV considers the average over 3 years and the maps only show the situation for one specific year.

Heatmaps with the evolution from 2000 of the mean (top) and the maximum (bottom) O<sub>3</sub> concentrations (93.2 percentile of the maximum daily 8-hour mean concentration, target value) at country level are shown in figure 19. In this way, the evolution along years of the average and maximum measured concentration levels can be seen for each country. Note that meteorological variability has a considerable impact on year-to-year changes in ambient air concentrations of air pollutants (EEA, 2020), especially for O<sub>3</sub> as higher ambient air temperature leads to enhanced photochemical reactions and O<sub>3</sub> formation. The last year (2024) is based on UTD data, while the previous years are based on officially reported validated data.

Figure 19: Evolution of mean (top) and maximum (bottom) O<sub>3</sub> concentrations (93.2 percentile of the maximum daily 8-hour mean concentration, related to the target value) per country from 2000



Note: It is important to note that the figure is not based on a consistent set of stations. The number, location and classification of the stations included may vary from year to year.



Figure 20: UTD Map of O<sub>3</sub> concentrations in 2024 - short-term WHO AQG level








Figure 21 shows the maps of the 99 percentile of the O<sub>3</sub> maximum daily 8-hour mean concentrations (equivalent to the short-term WHO AQG level) for the last four years. In this way, any significant change in the spatial distribution of the values above the set thresholds in the legends can be observed. Note that only the last year's map (2024) is based on UTD data, while the previous three years are based on officially reported validated data (CDR).

*Figure 21: Maps of O<sub>3</sub> concentrations (short-term WHO AQG level) for the last 4 years*




Heatmaps with the evolution from 2013 of the mean (top) and the maximum (bottom) 99 percentile of the O<sub>3</sub> maximum daily 8-hour mean concentrations at country level are shown in figure 22. In this way, the evolution along years of the average and maximum measured concentration levels can be seen for each country. Note that meteorological variability has a considerable impact on year-to-year changes in ambient air concentrations of air pollutants (EEA, 2020), and the last year (2024) is based on UTD data, while the previous years are based on officially reported validated data.

Figure 22: Evolution of mean (top) and maximum (bottom) 99 percentile of the O<sub>3</sub> maximum daily 8-hour mean concentrations per country from 2013



Note: It is important to note that the figure is not based on a consistent set of stations. The number, location and classification of the stations included may vary from year to year.


Figure 23: UTD Map of peak season O<sub>3</sub> concentrations in 2024



Note: Observed concentrations of O<sub>3</sub> in 2024. The map shows the average of the daily maximum 8-hour mean O<sub>3</sub> concentration in the six consecutive months with the highest six-month running-average O<sub>3</sub> concentration. The first colour category represents stations fulfilling the peak season O<sub>3</sub> AQG level. Only stations with more than 75 % of valid data have been included in the map.


Figure 24 shows the maps of the peak season O<sub>3</sub> concentrations (equivalent to the long-term WHO AQG level) for the last four years. In this way, any significant change in the spatial distribution of the values above the set thresholds in the legends can be observed. Note that only the last year's map (2024) is based on UTD data, while the previous three years are based on officially reported validated data (CDR).

Figure 24: Maps of peak season O<sub>3</sub> concentrations for the last 4 years



Heatmaps with the evolution from 2013 of the mean (top) and the maximum (bottom) peak season O<sub>3</sub> concentrations at country level are shown in figure 25. In this way, the evolution along years of the average and maximum measured concentration levels can be seen for each country. Note that meteorological variability has a considerable impact on year-to-year changes in ambient air concentrations of air pollutants (EEA, 2020), and the last year (2024) is based on UTD data, while the previous years are based on officially reported validated data.

Figure 25: Evolution of mean (top) and maximum (bottom) peak season O<sub>3</sub> concentrations per country from 2013



Note: It is important to note that the figure is not based on a consistent set of stations. The number, location and classification of the stations included may vary from year to year.

## 5 Status of nitrogen dioxide ambient air concentrations

The reporting countries shown in Figure 1 submitted NO<sub>2</sub> data from 2843 stations for the annual limit value, 2742 stations for the hourly limit value, and 2723 stations for the daily WHO AQG level.


6 of the countries in EU-27 and 1 other reporting countries (Figure 26) recorded concentrations above the annual limit value (40 µg/m<sup>3</sup>). This happened in 1 % of all the stations measuring NO<sub>2</sub>. On the contrary, 65 % of stations, located in 27 of the countries in EU-27 and 7 other reporting countries reported concentrations above the WHO AQG level of 10 µg/m<sup>3</sup>. Figure 26 shows the measured annual mean NO<sub>2</sub> concentrations.

96 % of all values above the annual limit value were observed at traffic stations. Furthermore, 100 % of the stations with concentrations above the annual limit value were located in urban or suburban areas.


Concentrations above the hourly limit value (200 µg/m<sup>3</sup>) were observed in 0.04 % (1 urban traffic station). It was observed in one countries. (Italy (one)).

Finally, concentrations above the daily NO<sub>2</sub> WHO AQG level (25 µg/m<sup>3</sup>) were registered in 72 % (1948 stations) of all the reporting stations in 27 of the countries in EU-27 and 6 other reporting countries (Figure 29).

Figure 26: UTD Map and boxplot of NO<sub>2</sub> concentrations in 2024



Note: Observed concentrations of NO<sub>2</sub> in 2024. The last two colour categories correspond to values above the EU annual limit value (40 µg/m<sup>3</sup>), while the first colour category indicates stations reporting values below the WHO AQG level for NO<sub>2</sub> (10 µg/m<sup>3</sup>). Only stations with more than 75 % of valid data have been included in the map.




Note: The graph is based on the annual mean concentration values. For each country, the number of stations considered for 2024 (in brackets) are given. The boxplot represents the lowest (bottom of the whisker), highest (top of the whisker) and average (black dot) annual mean values (in µg/m<sup>3</sup>). The rectangles mark the 25th and 75th percentiles. At 25 % of the stations, levels are below the 25th percentile; at 25 % of the stations, concentrations are above the 75th percentile. The limit value set by EU legislation is marked by the horizontal line. The WHO AQG level is marked by the lower dashed horizontal line. The graph should be read in relation to the above map, as a country's situation depends on the number of stations considered.



Figure 27 shows the maps of the observed NO<sub>2</sub> annual mean concentrations for the last four years. In this way, any significant change in the spatial distribution of the values above the set thresholds in the legends can be observed. Note that only the last year's map (2024) is based on UTD data, while the previous three years are based on officially reported validated data (CDR).

*Figure 27: Map of NO<sub>2</sub> concentrations (annual mean) for the last 4 years*




Maps for years before 2023 are different to the ones published in previous reports because the bands in the legend have been modified to accommodate the 2030 EU annual limit value (EU, 2024).

Heatmaps with the evolution from 2000 of the mean (top) and the maximum (bottom) NO<sub>2</sub> annual mean concentrations at country level are shown in figure 28. In this way, the evolution along years of the average and maximum measured concentration levels can be seen for each country. Note that meteorological variability has a considerable impact on year-to-year changes in ambient air concentrations of air pollutants (EEA, 2020), and the last year (2024) is based on UTD data, while the previous years are based on officially reported validated data.




Figure 28: Evolution of mean (top) and maximum (bottom) NO<sub>2</sub> annual mean concentrations (annual limit value) per country from 2000



Note: It is important to note that the figure is not based on a consistent set of stations. The number, location and classification of the stations included may vary from year to year.


Figure 29: UTD Map of NO<sub>2</sub> concentrations in 2024 - daily WHO AQG level



Note: Observed concentrations of NO<sub>2</sub> in 2024. The map shows the 99 percentile of the NO<sub>2</sub> daily mean concentrations, equivalent to 3–4 exceedance days per year, according to the definition of the daily WHO AQG level (25 µg/m<sup>3</sup>). The first colour category indicates stations with concentrations below this AQG level. Only stations with more than 75 % of valid data have been included in the map.


Figure 30 shows the maps of the 99 percentile of NO<sub>2</sub> daily mean concentrations (equivalent to the WHO AQG level for NO<sub>2</sub> daily mean level) for the last four years. In this way, any significant change in the spatial distribution of the values above the set thresholds in the legends can be observed. Note that only the last year's map (2024) is based on UTD data, while the previous three years are based on officially reported validated data (CDR).

*Figure 30: Maps of NO<sub>2</sub> concentrations (daily WHO AQG level) for the last 4 years*



Heatmaps with the evolution from 2013 of the mean (top) and the maximum (bottom) 99 percentile of NO<sub>2</sub> daily mean concentrations at country level are shown in figure 31. In this way, the evolution along years of the average and maximum measured concentration levels can be seen for each country. Note that meteorological variability has a considerable impact on year-to-year changes in ambient air concentrations of air pollutants (EEA, 2020), and the last year (2024) is based on UTD data, while the previous years are based on officially reported validated data.

Figure 31: Evolution of mean (top) and maximum (bottom) 99 percentile of NO<sub>2</sub> daily mean concentrations (daily WHO AQG level) per country from 2013



Note: It is important to note that the figure is not based on a consistent set of stations. The number, location and classification of the stations included may vary from year to year.

## 6 Status of sulphur dioxide ambient air concentrations

The reporting countries shown in Figure 1 reported measurements of SO<sub>2</sub> from 1159 stations for the hourly limit value and 1144 stations for the daily limit value.

7 stations <sup>(5)</sup> registered concentrations above the hourly limit value (350 µg/m<sup>3</sup>); and 5 stations <sup>(6)</sup> registered concentrations above the daily limit of 125 µg/m<sup>3</sup> for SO<sub>2</sub> (Figure 32).

On the contrary, 32 (3 %) of all the stations reporting SO<sub>2</sub> levels, located in 12 reporting countries <sup>(7)</sup>, measured SO<sub>2</sub> concentrations above the WHO AQG level of 40 µg/m<sup>3</sup> for daily mean concentrations <sup>(8)</sup>.

---


<sup>5</sup>Italy (six) and Iceland (one)

<sup>6</sup>Italy (two), Greece (one), Iceland (one) and North Macedonia (one).

<sup>7</sup>All reporting countries except Andorra, Austria, Belgium, Croatia, Cyprus, Denmark, Estonia, Finland, Ireland, Kosovo, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Portugal, Romania, Slovenia and Switzerland.

<sup>8</sup>Although the WHO AQG level for daily means refers to the percentile 99 (3-4 exceedance days), here we have used the percentile 99.18 (3 exceedance days), so the daily WHO AQG level can be directly compared with the EU daily LV.


Figure 32: Map of SO<sub>2</sub> daily concentrations in 2024



Note: Observed concentrations of SO<sub>2</sub> in 2024. The map shows the percentile 99.18 of SO<sub>2</sub> daily means, indicating 3 exceedance days. It relates to the EU daily limit value (125 µg/m<sup>3</sup>) and to the WHO daily AQG level (40 µg/m<sup>3</sup>). Only stations with more than 75 % of valid data have been included in the map.

Figure 33 shows the maps of the observed SO<sub>2</sub> daily mean concentrations for the last four years. In this way, any significant change in the spatial distribution of the values above the set thresholds in the legends can be observed. Note that only the last year's map (2024) is based on UTD data, while the previous three years are based on officially reported validated data (CDR).


*Figure 33: Maps of SO<sub>2</sub> concentrations (daily mean) for the last 4 years*



Heatmaps with the evolution from 2000 of the mean (top) and the maximum (bottom) SO<sub>2</sub> daily mean concentrations at country level are shown in figure 34. In this way, the evolution along years of the average and maximum measured concentration levels can be seen for each country. Note that meteorological variability has a considerable impact on year-to-year changes in ambient air concentrations of air pollutants (EEA, 2020), and the last year (2024) is based on UTD data, while the previous years are based on officially reported validated data.



Figure 34: Evolution of mean (top) and maximum (bottom) SO<sub>2</sub> 99.18 percentile of daily mean concentrations (EU LV (125 µg/m<sup>3</sup>) and WHO AQG level (40 µg/m<sup>3</sup>)) per country from 2000



Note: It is important to note that the figure is not based on a consistent set of stations. The number, location and classification of the stations included may vary from year to year.



## 7 Abbreviations, units and symbols

$\mu\text{g}/\text{m}^3$ : microgram(s) per cubic metre

AAQD: Ambient Air Quality Directives

AQG: Air quality guideline

CDR: Central data repository

EEA: European Environment Agency

ETC HE: European Topic Centre on Human health and the Environment

EU: European Union

LV: limit value

$\text{NO}_2$ : Nitrogen dioxide

$\text{O}_3$ : Ozone

PM: Particulate matter

$\text{PM}_{2.5}$ : Particulate matter with a diameter of 2.5  $\mu\text{m}$  or less

$\text{PM}_{10}$ : Particulate matter with a diameter of 10  $\mu\text{m}$  or less

RL: Reference level

$\text{SO}_2$ : Sulphur dioxide

TV: target value

UTD: up-to-date

WHO: World Health Organization

## 8 Annex 1

Data included in this report was received by 19 February 2025 from the reporting countries. By that date the number of stations by country aggregation reporting each pollutant is summarized in Table 3. Data from stations that do not fulfil the criteria from Box 1.1 are excluded from this report.

*Table 3: Reporting status of 2024 air quality data by 19 February 2025*

Countries	PM10	PM2.5	O3	NO2	SO2
EU27	2402	1519	1894	2735	1114
EEA32	2489	1580	1936	2813	1138
Total	2514	1605	1973	2843	1163

Data not included in this report is summarized in Table 4:

*Table 4: Reporting outliers of 2024 air quality data by 19 February 2025*

Country	Station Eol Code	Pollutant	Aggregation(*)	Year	Value	Units	Data Coverage
Italy	IT1486A	PM10	P1Y-P1D- per90.41	2024	973	ug/m3	83
Italy	IT1486A	PM10	P1Y-P1D- per99	2024	3065	ug/m3	83
Italy	IT2132A	SO2	P1Y	2024	101498	ug/m3	95
Italy	IT2132A	SO2	P1Y-day-max- per99.18	2024	269846	ug/m3	98
Italy	IT2133A	PM10	P1Y	2024	1485	ug/m3	95
Italy	IT2148A	PM10	P1Y-P1D- per99	2024	591	ug/m3	94
Italy	IT2148A	PM2.5	P1Y-P1D- per99	2024	985	ug/m3	93
Italy	IT2151A	PM2.5	P1Y-P1D- per99	2024	985	ug/m3	93
Italy	IT2261A	PM10	P1Y	2024	1418	ug/m3	94

Table 4: Reporting outliers of 2024 air quality data by 19 February 2025 (continued)

Country	Station Eol Code	Pollutant	Aggregation(*)	Year	Value	Units	Data Coverage
Italy	IT2261A	PM10	P1Y-P1D- per99	2024	15628	ug/m3	95
Italy	IT1486A	PM10	P1Y-P1D- per95.07	2024	1885		83
Italy	IT2148A	PM2.5	P1Y-P1D- per95.07	2024	985		93
Italy	IT2151A	PM2.5	P1Y-P1D- per95.07	2024	646		93

(\*) <https://dd.eionet.europa.eu/vocabulary/aq/aggregationprocess/view>

Table 5 summarizes the number of sampling points per country with air quality levels above specific air quality objectives summarized through out this report. Sampling points that do not fulfil the criteria from Box 1.1 are excluded.

*Table 5: Number of sampling points above air quality levels/objectives per reporting country*

Levels/Objectives	EU Member States																																	EU Institutions			EU Candidate Countries			EU Potential Candidate Countries			EU Neighbouring Countries			EU Overseas Countries and Territories		
	Andorra	Austria	Belgium	Bulgaria	Croatia	Cyprus	Czechia	Denmark	Estonia	Finland	France	Germany	Greece	Hungary	Iceland	Ireland	Italy	Kosovo	Latvia	Lithuania	Luxembourg	Malta	Netherlands	North Macedonia	Norway	Poland	Portugal	Romania	Serbia	Slovakia	Slovenia	Spain	Sweden	Switzerland														
PM <sub>10</sub> daily LV (50 µg/m <sup>3</sup> )	0	0	0	1	2	1	0	0	0	0	3	0	5	0	0	0	87	0	0	0	0	3	0	11	1	10	0	6	0	3	2	38	1	0														
PM <sub>10</sub> daily 2030 LV (45 µg/m <sup>3</sup> )	0	3	1	13	5	2	14	0	1	2	14	1	14	7	0	0	230	3	2	3	0	0	0	11	13	88	3	10	0	13	3	100	9	0														
PM <sub>10</sub> daily WHO AQG level (45 µg/m <sup>3</sup> )	1	77	9	20	9	2	65	0	1	11	64	85	20	22	2	2	414	7	7	13	0	5	2	11	36	163	51	36	0	41	18	244	30	0														
PM <sub>10</sub> annual LV (40 µg/m <sup>3</sup> )	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	0	0	0	0	0	0	6	1	0	0	0	0	0	0	4	0	0														
PM <sub>10</sub> annual WHO AQG level (15 µg/m <sup>3</sup> )	0	54	44	22	6	2	57	1	1	3	98	75	20	20	1	2	397	7	7	10	2	5	26	11	20	163	43	41	0	46	17	252	19	0														
PM <sub>10</sub> annual 2030 LV (20 µg/m <sup>3</sup> )	0	10	4	16	4	2	22	0	0	0	25	8	17	11	0	0	293	1	2	5	0	5	1	11	5	122	15	23	0	25	12	144	2	0														
PM <sub>2.5</sub> annual LV (25 µg/m <sup>3</sup> )	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	6	0	0	0	0	0	0	0	0	0	0														
PM <sub>2.5</sub> annual WHO AQG level (5 µg/m <sup>3</sup> )	1	53	63	3	10	1	48	0	0	3	213	312	10	10	0	16	224	12	5	6	3	5	24	11	38	89	11	12	1	47	18	214	15	7														
PM <sub>2.5</sub> daily WHO AQG level (15 µg/m <sup>3</sup> )	1	53	71	3	10	1	48	0	3	9	219	319	10	10	1	15	222	12	5	5	3	5	24	11	43	89	16	12	0	47	18	222	20	7														
O <sub>3</sub> max daily 8h mean TV (120 µg/m <sup>3</sup> )	0	2	0	2	3	3	4	0	0	0	2	2	10	5	0	0	127	1	0	0	0	0	0	3	0	7	2	3	0	4	4	59	0	2														
O <sub>3</sub> long-term objective (120 µg/m <sup>3</sup> )	1	87	33	14	13	3	54	7	5	3	211	259	16	16	0	1	239	8	2	10	4	3	39	7	4	86	37	12	0	19	11	275	13	26														
O <sub>3</sub> max daily 8h mean 2030 LTO (100 µg/m <sup>3</sup> )	0	3	0	4	7	3	15	0	0	0	7	12	12	7	0	0	152	3	0	0	0	0	0	3	0	19	4	3	0	6	5	71	0	4														
O <sub>3</sub> short-term WHO AQG level (100 µg/m <sup>3</sup> )	1	106	33	15	14	3	54	8	9	13	265	275	17	16	0	1	261	8	5	13	5	3	40	10	10	95	42	16	0	23	11	373	17	30														
O <sub>3</sub> peak season WHO AQG level (60 µg/m <sup>3</sup> )	1	93	33	19	11	3	55	8	9	14	276	275	18	16	0	6	262	6	6	12	5	4	40	9	11	95	35	17	8	22	11	409	18	30														
NO <sub>2</sub> annual LV (40 µg/m <sup>3</sup> )	0	0	0	0	0	0	0	0	0	0	4	1	4	0	0	0	13	0	0	0	0	0	0	1	0	2	2	0	0	0	0	0	0	0														
NO <sub>2</sub> annual WHO AQG level (10 µg/m <sup>3</sup> )	1	100	59	18	10	2	36	3	1	11	188	356	14	17	3	8	408	6	4	9	6	4	52	14	33	91	38	19	1	21	7	275	24	23														
NO <sub>2</sub> hourly LV (200 µg/m <sup>3</sup> )	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0														
NO <sub>2</sub> hourly 2030 LV (200 µg/m <sup>3</sup> )	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	1	14	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0														
NO <sub>2</sub> daily WHO AQG level (25 µg/m <sup>3</sup> )	1	117	65	19	12	2	45	3	3	21	225	277	14	16	3	10	419	6	5	10	6	4	62	16	37	90	39	20	0	25	8	316	27	25														
NO <sub>2</sub> daily 2030 LV (50 µg/m <sup>3</sup> )	0	2	1	5	1	0	1	0	0	0	9	2	8	1	0	2	64	0	0	1	0	0	1	2	6	4	3	4	0	1	1	19	1	0														
SO <sub>2</sub> hourly LV (350 µg/m <sup>3</sup> )	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0														
SO <sub>2</sub> hourly LV 2030 (350 µg/m <sup>3</sup> )	0	1	0	2	0	0	4	0	0	0	1	1	0	0	2	0	12	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0														
SO <sub>2</sub> daily LV (125 µg/m <sup>3</sup> )	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	2	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0														
SO <sub>2</sub> daily WHO AQG level (40 µg/m <sup>3</sup> )	0	0	0	3	0	0	4	0	0	0	1	3	1	2	4	0	5	0	0	0	0	0	0	2	0	1	0	0	0	1	0	5	0	0														
CO daily LV (10 mg/m <sup>3</sup> )	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	1	9	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0														
CO daily WHO AQG level (4 mg/m <sup>3</sup> )	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0														

Table 5: Number of sampling points above air quality levels/objectives per reporting country (continued)

Levels/Objectives	Member States																																	
	Andorra	Austria	Belgium	Bulgaria	Croatia	Cyprus	Czechia	Denmark	Estonia	Finland	France	Germany	Greece	Hungary	Iceland	Ireland	Italy	Kosovo	Latvia	Lithuania	Luxembourg	Malta	Netherlands	North Macedonia	Norway	Poland	Portugal	Romania	Serbia	Slovakia	Slovenia	Spain	Sweden	Switzerland
C <sub>6</sub> H <sub>6</sub> annual LV (5 µg/m <sup>3</sup> )	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0
C <sub>6</sub> H <sub>6</sub> annual WHO RL (1.7 µg/m <sup>3</sup> )	0	0	2	3	1	0	0	0	0	0	0	0	0	1	0	0	14	0	0	0	0	0	0	0	0	2	0	9	0	0	0	2	0	0
C <sub>6</sub> H <sub>6</sub> annual 2030 LV (3.4 µg/m <sup>3</sup> )	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0

## 9 Annex 2 - Air quality status in relation to the new standards in the revised AAQD (EU) 2024/2081

On 20 November 2024, the revised Directive (EU) 2024/2881 (EU, 2024) on ambient air quality and cleaner air for Europe was published and it entered into force on 10 December 2024. This Directive amends and recasts Directives 2004/107/EC (EU, 2004) and 2008/50/EC (EU, 2008). Among other changes, it has introduced new air quality (AQ) standards (or revised the ones in the 2004 and 2008 Directives) setting the attainment date at 1 January 2030. The revised and new standards for the protection of human health are summarised in table 6).

*Table 6: Air quality standards for the protection of health, as given in the revised EU Ambient Air Quality Directive (EU) 2024/2881*

Pollutant	Averaging period	Legal nature and concentration	Comments
PM <sub>10</sub>	1 day	Limit value: 45 µg/m <sup>3</sup>	Not to be exceeded more than 18 times per calendar year
		Alert threshold: 90 µg/m <sup>3</sup>	To be measured as a daily average over 3 consecutive days or less at locations representative of air quality over at least 100 km <sup>2</sup> or an entire zone, whichever is the smaller
		Information threshold: 90 µg/m <sup>3</sup>	To be measured over 1 day at locations representative of air quality over at least 100 km <sup>2</sup> or an entire zone, whichever is the smaller
	Calendar year	Limit value: 20 µg/m <sup>3</sup>	
PM <sub>2.5</sub>	1 day	Limit value: 25 µg/m <sup>3</sup>	Not to be exceeded more than 18 times per calendar year
		Alert threshold: 50 µg/m <sup>3</sup>	To be measured as a daily average over 3 consecutive days or less at locations representative of air quality over at least 100 km <sup>2</sup> or an entire zone, whichever is the smaller
		Information threshold: 50 µg/m <sup>3</sup>	To be measured over 1 day at locations representative of air quality over at least 100 km <sup>2</sup> or an entire zone, whichever is the smaller
	Calendar year	Limit value: 10 µg/m <sup>3</sup>	
		Average exposure reduction obligation, 10-25% reduction of the Average Exposure Indicator (AEI) in 2020	The percentage reduction depends on the initial AEI in 2020 <sup>(a)</sup>
O <sub>3</sub>	1 hour	Alert threshold: 240 µg/m <sup>3</sup>	
		Information threshold: 180 µg/m <sup>3</sup>	
	Maximum daily 8-hour mean <sup>(b)</sup>	Target value: 120 µg/m <sup>3</sup>	Not to be exceeded on more than 18 days per calendar year averaged over 3 years <sup>(c)</sup>
		Average exposure concentration objective, AIE: 5 µg/m <sup>3</sup>	

*Table 6: Air quality standards for the protection of health, as given in the revised EU Ambient Air Quality Directive (EU) 2024/2881 (continued)*

Pollutant	Averaging period	Legal nature and concentration	Comments
NO <sub>2</sub>	Maximum daily 8-hour mean within a calendar year	Long term objective: 100 µg/m <sup>3</sup>	Not to be exceeded more than 3 days per calendar year (99th percentile)
	1 hour	Limit value: 200 µg/m <sup>3</sup>	Not to be exceeded more than 3 times per calendar year
		Alert threshold: 200 µg/m <sup>3</sup>	To be measured as a daily average over 3 consecutive days or less at locations representative of air quality over at least 100 km <sup>2</sup> or an entire zone, whichever is the smaller
		Information threshold: 150 µg/m <sup>3</sup>	To be measured over 1 day at locations representative of air quality over at least 100 km <sup>2</sup> or an entire zone, whichever is the smaller
	1 day	Limit value: 50 µg/m <sup>3</sup>	Not to be exceeded more than 18 times per calendar year
	Calendar year	Limit value: 20 µg/m <sup>3</sup>	
		Average exposure reduction obligation, 15-25% reduction of the Average Exposure Indicator (AEI) in 2020	The percentage reduction depends on the initial AEI in 2020 <sup>(a)</sup>
		Average exposure concentration objective, AIE: 10 µg/m <sup>3</sup>	
BaP	Calendar year	Limit value: 1,0 ng/m <sup>3</sup>	
SO <sub>2</sub>	1 hour	Limit value: 350 µg/m <sup>3</sup>	Not to be exceeded more than 3 times per calendar year
		Alert threshold: 350 µg/m <sup>3</sup>	To be measured as a daily average over 3 consecutive days or less at locations representative of air quality over at least 100 km <sup>2</sup> or an entire zone, whichever is the smaller
		Information threshold: 275 µg/m <sup>3</sup>	To be measured over 1 day at locations representative of air quality over at least 100 km <sup>2</sup> or an entire zone, whichever is the smaller
	1 day	Limit value: 50 µg/m <sup>3</sup>	Not to be exceeded more than 18 times per calendar year
CO	Maximum daily 8-hour mean <sup>(b)</sup>	Limit value: 10 mg/m <sup>3</sup>	
	1 day	Limit value: 4 mg/m <sup>3</sup>	Not to be exceeded more than 18 times per calendar year
C <sub>6</sub> H <sub>6</sub>	Calendar year	Limit value: 3.4 µg/m <sup>3</sup>	
Pb	Calendar year	Limit value: 0.5 µg/m <sup>3</sup>	
As	Calendar year	Limit value: 6,0 ng/m <sup>3</sup>	
Cd	Calendar year	Limit value: 5,0 ng/m <sup>3</sup>	
Ni	Calendar year	Limit value: 20 ng/m <sup>3</sup>	

Table 6: Air quality standards for the protection of health, as given in the revised EU Ambient Air Quality Directive (EU) 2024/2881 (continued)

Pollutant	Averaging period	Legal nature and concentration	Comments
<p><b>Notes:</b></p> <p><sup>a</sup> AEI: based upon measurements at all sampling points in urban background locations in average exposure territorial units throughout the territory of a Member State, assessed as a 3-year running annual mean.</p> <p><sup>b</sup> The maximum daily 8-hour mean concentration shall be selected by examining 8-hour running averages, calculated from hourly data and updated each hour. Each 8-hour average so calculated shall be assigned to the day on which it ends, i.e. the first calculation period for any 1 day shall be the period from 17:00 on the previous day to 1:00 on that day; the last calculation period for any 1 day shall be the period from 16:00 to 24:00 on that day.</p> <p><sup>c</sup> If the 3-year average cannot be determined on the basis of a full and consecutive set of annual data, the minimum annual data required for checking compliance with the ozone target value shall be valid data for 1 year.</p>			

This annex analyses the situation in year 2024 with respect to some of the new and/or revised AQ standards <sup>(9)</sup> using the same air quality data as those in the main section. It provides:

- a European overview of the 2024 monitoring stations reported, and of their concentrations in relation to some EU legal standards defined in the Directive (EU) 2024/2881 (EU, 2024);
- a map with the 2024 concentrations at station level for each pollutant relevant to each AQ standard;
- a boxplot graph summarizing for each country the range of concentrations (highlighting the lowest, highest, average and the 25 and 75 percentiles) for PM<sub>10</sub>, PM<sub>2.5</sub>, NO<sub>2</sub> and O<sub>3</sub>.

It should be noted that this assessment is only intended to shed light on the current situation in relation to the 2030 AQ standards, and the ‘distance to target’, i.e. the scale of the challenge to meet those standards by 2030. This analysis may also help to identify air quality zones where air quality roadmaps may need to be established. Roadmaps are air quality plans that must be implemented where, from 2026 to 2029, levels of pollutants are foreseen to be above the 2030 limit and target values.

<sup>9</sup>Only for the limit values and O3 target value and long-term objective




## 9.1 Status of PM<sub>10</sub> concentrations


Twenty-two countries in EU-27, and three other reporting countries reported PM<sub>10</sub> concentrations above the 2030 EU daily limit value of 45 µg/m<sup>3</sup>, more than 18 days over the calendar year (Figure 35). This applied to 23 % (565) of reporting stations.

Concentrations above the 2030 PM<sub>10</sub> annual limit value (20 µg/m<sup>3</sup>) were monitored in 31 % (785 stations) of all the reporting stations, located in 22 countries in EU-27, and 3 other reporting countries (Figure 36).

Figure 35: UTD Map and boxplot of PM<sub>10</sub> concentrations in 2024 - 2030 daily limit value




Note: The figure shows observed concentrations of PM<sub>10</sub> in 2024. The possibility of subtracting contributions to the measured concentrations from natural sources and winter road sanding/salting has not been considered. The map shows the 95.07 percentile of the PM<sub>10</sub> daily mean concentrations, representing the 19th highest value in a complete series. It is related to the 2030 PM<sub>10</sub> daily limit value, allowing 18 exceedances of the 45 µg/m<sup>3</sup> threshold over 1 year. The last two colour categories indicate stations with concentrations above this daily limit value. Only stations with more than 75 % of valid data have been included in the map.




Note: The graph is based on the 95.07 percentile of daily mean concentration values corresponding to the 19th highest daily mean in complete time series. For each country, the number of stations considered for 2024 (in brackets) are given. The boxplot represents the lowest (bottom of the whisker), highest (top of the whisker) and average (black dot) 95.07 percentile values (in µg/m<sup>3</sup>). The rectangles mark the 25th and 75th percentiles. At 25% of the stations, levels are below the 25th percentile; at 25 % of the stations, concentrations are above the 75th percentile. The daily limit value set by EU legislation for 2030 is marked by the horizontal line. The graph should be read in relation to the above map, as a country's situation depends on the number of stations considered.

Figure 36: UTD Map and Boxplot of  $PM_{10}$  concentrations in 2024 - 2030 annual limit value



Note: Observed concentrations of  $PM_{10}$  in 2024. The possibility of subtracting contributions to the measured concentrations from natural sources and winter road sanding/salting has not been considered. The last three colour categories indicate stations reporting concentrations above the 2030 EU annual limit value ( $20 \mu g/m^3$ ). Only stations with more than 75 % of valid data, have been included in the map.




Note: The graph is based on annual mean concentration values. For each country, the number of stations considered for 2024 (in brackets) are given. The boxplot represents the lowest (bottom of the whisker), highest (top of the whisker) and average (black dot) annual mean values (in  $\mu g/m^3$ ). The rectangles mark the 25th and 75th percentiles. At 25 % of the stations, levels are below the 25th percentile; at 25 % of the stations, concentrations are above the 75th percentile. The annual limit value set by EU legislation for 2030 is marked by the continuous horizontal line. The graph should be read in relation to the above map, as a country's situation depends on the number of stations considered.

## 9.2 Status of PM<sub>2.5</sub> concentrations


Seventeen countries in EU-27, and three other reporting countries reported PM<sub>2.5</sub> concentrations above the 2030 EU daily limit value of 25 µg/m<sup>3</sup>, more than 18 days over the calendar year (Figure 37). This was the case for 27 % (410) of reporting stations (Figure 37).

The PM<sub>2.5</sub> concentrations were higher than the 2030 EU annual limit value (10 µg/m<sup>3</sup>) in twenty countries in EU-27 and four other reporting countries (Figure 38). These concentrations above the limit value were registered in 34 % (542 stations) of all the reporting stations.

Figure 37: UTD Map and boxplot of PM<sub>2.5</sub> concentrations in 2024 - 2030 daily limit value




Note: The figure shows observed concentrations of PM<sub>2.5</sub> in 2024. The possibility of subtracting contributions to the measured concentrations from natural sources has not been considered. The map shows the 95.07 percentile of the PM<sub>2.5</sub> daily mean concentrations, representing the 19th highest value in a complete series. It is related to the 2030 PM<sub>2.5</sub> daily limit value, allowing 18 exceedances of the 25 µg/m<sup>3</sup> threshold over 1 year. The last two colour categories indicate stations with concentrations above this daily limit value. Only stations with more than 75 % of valid data have been included in the map.




Note: The graph is based on the 95.07 percentile of daily mean concentration values corresponding to the 19th highest daily mean in complete time series. For each country, the number of stations considered for 2024 (in brackets) are given. The boxplot represents the lowest (bottom of the whisker), highest (top of the whisker) and average (black dot) 95.07 percentile values (in µg/m<sup>3</sup>). The rectangles mark the 25th and 75th percentiles. At 25% of the stations, levels are below the 25th percentile; at 25 % of the stations, concentrations are above the 75th percentile. The daily limit value set by EU legislation for 2030 is marked by the horizontal line. The graph should be read in relation to the above map, as a country's situation depends on the number of stations considered.

Figure 38: UTD Map and Boxplot of PM<sub>2.5</sub> concentrations in 2024 - 2030 annual limit value



Note: Observed concentrations of PM<sub>2.5</sub> in 2024. The possibility of subtracting contributions to the measured concentrations from natural sources has not been considered. The last three colour categories indicate stations reporting concentrations above the 2030 EU annual limit value (10 µg/m<sup>3</sup>). Only stations with more than 75 % of valid data, have been included in the map.




Note: The graph is based on annual mean concentration values. For each country, the number of stations considered for 2024 (in brackets) are given. The boxplot represents the lowest (bottom of the whisker), highest (top of the whisker) and average (black dot) annual mean values (in µg/m<sup>3</sup>). The rectangles mark the 25th and 75th percentiles. At 25 % of the stations, levels are below the 25th percentile; at 25 % of the stations, concentrations are above the 75th percentile. The annual limit value set by EU legislation for 2030 is marked by the continuous horizontal line. The graph should be read in relation to the above map, as a country's situation depends on the number of stations considered.

### 9.3 Status of O<sub>3</sub> ambient air concentrations


16 countries in EU-27 and 3 other reporting countries reported concentrations above the 2030 O<sub>3</sub> target value threshold (120 µg/m<sup>3</sup>) more than 18 times this year (Figure 39). In total, 18 % (340 stations) of all stations reporting O<sub>3</sub> showed concentrations above the 2030 target value threshold for the protection of human health.

Since the 2030 long-term objective aligns with the short-term WHO AQG (100 µg/m<sup>3</sup> as percentile 99), please refer to the main text and Figure 20 for an overview of the current situation in relation to the 2030 long-term objective.

Figure 39: UTD Map and boxplot of O<sub>3</sub> concentrations in 2024 - 2030 target value threshold.



Note: Observed concentrations of O<sub>3</sub> in 2024. The map shows the 95.07 percentile of the O<sub>3</sub> maximum daily 8-hour mean, representing the 19th highest value in a complete series. It is related to the 2030 O<sub>3</sub> target value. At sites marked with the last two colour categories, the 19th highest daily O<sub>3</sub> concentrations were above the 120 µg/m<sup>3</sup> threshold, implying values above the 2030 target value threshold. Please note that the legal definition of the target value considers not only 1 year but the average over 3 years.



Note: The graph is based, for each country, on the 95.07 percentile of the maximum daily 8-hour mean concentration values, corresponding to the 19th highest daily maximum of the running 8-hour mean in a complete time series. For each country, the number of stations considered for 2024 (in brackets) are given. The boxplot represents the lowest (bottom of the whisker), highest (top of the whisker) and average (black dot) values (in µg/m<sup>3</sup>). The rectangles mark the 25th and 75th percentiles. At 25 % of the stations, levels are below the 25th percentile; at 25% of the stations, concentrations are above the 75th percentile. The target value threshold set for 2030 by the EU legislation is marked by the horizontal line. Please note that the legal definition of the target value considers not only 1 year but the average over 3 years. The graph should be read in relation to the above map, as a country's situation depends on the number of stations considered.



#### 9.4 Status of NO<sub>2</sub> ambient air concentrations


23 of the countries in EU-27 and 4 other reporting countries (Figure 40) recorded concentrations above the 2030 EU annual limit value (20 µg/m<sup>3</sup>). This happened in 22 % (637 stations) of all the stations measuring NO<sub>2</sub>.

77 % (489 stations) of all values above the 2030 EU annual limit value were observed at traffic stations.


Concentrations above the 2030 EU NO<sub>2</sub> daily limit value (50 µg/m<sup>3</sup>, not to be exceeded more than 18 days per year) were reported in 5 % (139 stations) of all the reporting stations in 20 of the countries in EU-27 and 2 other reporting countries (Figure 41).

Finally, concentrations above the hourly limit value (200 µg/m<sup>3</sup>, not to be exceeded more than 3 hours per year) were observed in 0.8 % (23 stations) of all reporting stations, mostly at urban traffic stations. They were observed in eight countries (number stations): Italy (fourteen), Latvia (two), Spain (two), Belgium (one), Germany (one), Ireland (one), Slovakia (one) and Sweden (one).

Figure 40: UTD Map and Boxplot of NO<sub>2</sub> concentrations in 2024 - 2030 annual limit value




Note: Observed concentrations of NO<sub>2</sub> in 2024. The last three colour categories indicate stations reporting concentrations above the 2030 EU annual limit value (20 µg/m<sup>3</sup>). Only stations with more than 75 % of valid data have been included in the map.




Note: The graph is based on annual mean concentration values. For each country, the number of stations considered for 2024 (in brackets) are given.

The boxplot represents the lowest (bottom of the whisker), highest (top of the whisker) and average (black dot) annual mean values (in µg/m<sup>3</sup>). The rectangles mark the 25th and 75th percentiles. At 25 % of the stations, levels are below the 25th percentile; at 25 % of the stations, concentrations are above the 75th percentile. The annual limit value set by EU legislation for 2030 is marked by the continuous horizontal line. The graph should be read in relation to the above map, as a country's situation depends on the number of stations considered.

Figure 41: UTD Map and boxplot of NO<sub>2</sub> concentrations in 2024 - 2030 daily limit value



Note: The figure shows observed concentrations of NO<sub>2</sub> in 2024. The map shows the 95.07 percentile of the NO<sub>2</sub> daily mean concentrations, representing the 19th highest value in a complete series. It is related to the 2030 NO<sub>2</sub> daily limit value, allowing 18 exceedances of the 50 µg/m<sup>3</sup> threshold over 1 year. The last two colour categories indicate stations with concentrations above this daily limit value. Only stations with more than 75 % of valid data have been included in the map.



Note: The graph is based on the 95.07 percentile of daily mean concentration values corresponding to the 19th highest daily mean in complete series. For each country, the number of stations considered for 2024 (in brackets) are given. The boxplot represents the lowest (bottom of the whisker) and highest (top of the whisker) and average (black dot) 95.07 percentile values (in µg/m<sup>3</sup>). The rectangles mark the 25th and 75th percentiles. At 25% of the stations, levels are below the 25th percentile; at 25 % of the stations, concentrations are above the 75th percentile. The daily limit value EU legislation for 2030 is marked by the horizontal line. The graph should be read in relation to the above map, as a country's situation depends on the number of stations considered.


## 9.5 Status of SO<sub>2</sub> concentrations

3 of the countries in EU-27 and 1 other reporting countries (Figure 42) recorded concentrations above the 2030 EU annual limit value (20 µg/m<sup>3</sup>). This happened in 0.4 % of all the stations measuring SO<sub>2</sub>.


Concentrations above the 2030 EU SO<sub>2</sub> daily limit value (50 µg/m<sup>3</sup>, not to be exceeded more than 18 days per year) were registered in 1 % (6 stations) of all the reporting stations in 3 of the countries in EU-27 and 2 other reporting countries (Figure 43).

Finally, concentrations above the hourly limit value (350 µg/m<sup>3</sup> not to be exceeded in more than 3 occasions) were observed in 2.1 % (24 stations) of all reporting stations. They were observed in eight countries (number stations): Italy (twelve), Czechia (four), Bulgaria (two), Iceland (two), Austria (one), France (one), Germany (one) and North Macedonia (one).

Figure 42: UTD Map and Boxplot of SO<sub>2</sub> concentrations in 2024 - 2030 annual limit value




Note: Observed concentrations of SO<sub>2</sub> in 2024. The last two colour categories indicate stations reporting concentrations above the 2030 EU annual limit value (20 µg/m<sup>3</sup>). Only stations with more than 75 % of valid data have been included in the map.



Note: The graph is based on annual mean concentration values. For each country, the number of stations considered for 2024 (in brackets) are given. The boxplot represents the lowest (bottom of the whisker), highest (top of the whisker) and average (black dot) annual mean values (in µg/m<sup>3</sup>). The rectangles mark the 25th and 75th percentiles. At 25 % of the stations, levels are below the 25th percentile; at 25 % of the stations, concentrations are above the 75th percentile. The annual limit value set by EU legislation for 2030 is marked by the continuous horizontal line. The graph should be read in relation to the above map, as a country's situation depends on the number of stations considered.


Figure 43: UTD Map and boxplot of SO<sub>2</sub> concentrations in 2024 - 2030 daily limit value



Note: The figure shows observed concentrations of SO<sub>2</sub> in 2024. The map shows the 95.07 percentile of the SO<sub>2</sub> daily mean concentrations, representing the 19th highest value in a complete series. It is related to the 2030

SO<sub>2</sub> daily limit value, allowing 18 exceedances of the 50 µg/m<sup>3</sup> threshold over 1 year. The last two colour categories indicate stations with concentrations above this daily limit value. Only stations with more than 75 % of valid data have been included in the map.

**SO<sub>2</sub> concentrations in 2024 in relation to the 2030 standard daily limit value and number of stations considered for each country**



Note: The graph is based on the 95.07 percentile of daily mean concentration values corresponding to the 19th highest daily mean in complete series. For each country, the number of stations considered for 2024 (in brackets) are given. The boxplot represents the lowest (bottom of the whisker) and highest (top of the whisker) and average (black dot) 95.07 percentile values (in µg/m<sup>3</sup>). The rectangles mark the 25th and 75th percentiles. At 25% of the stations, levels are below the 25th percentile; at 25 % of the stations, concentrations are above the 75th percentile. The daily limit value set by EU legislation for 2030 is marked by the horizontal line. The graph should be read in relation to the above map, as a country's situation depends on the number of stations considered.

## References

- EEA (2020). Air quality in Europe–2020 report. *EEA Report No 9/2020*, <https://www.eea.europa.eu/publications/air-quality-in-europe-2020>.
- EU (2004). Directive 2004/107/EC of the European Parliament and of the Council of 15 December 2004 relating to arsenic, cadmium, mercury, nickel and polycyclic aromatic hydrocarbons in ambient air. *OJ L 23*, 26.1.2005, p. 3–16.
- EU (2008). Directive 2008/50/EC of the European Parliament and of the Council of 21 May 2008 on ambient air quality and cleaner air for Europe. *OJ L 152*, 11.6.2008, pp. 1–44.
- EU (2011). Commission Implementing Decision No 2011/850/EU of 12 December 2011 laying down rules for Directives 2004/107/EC and 2008/50/EC of the European Parliament and of the Council as regards the reciprocal exchange of information and reporting on ambient air quality. *OJ L 335*, 17.12.2011, pp. 86–106.
- EU (2024). Directive (EU) 2024/2881 of the European Parliament and of the Council of 23 October 2024 on ambient air quality and cleaner air for Europe.
- WHO (2000). Air quality guidelines for Europe, *World Health Organization, Regional Office for Europe, Copenhagen*.
- WHO (2006). Air quality guidelines: Global update 2005 — Particulate matter, ozone, nitrogen dioxide and sulphur dioxide, *World Health Organization, Regional Office for Europe, Copenhagen*.
- WHO (2021). WHO global air quality guidelines. Particulate matter (PM<sub>2.5</sub> and PM<sub>10</sub>), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide. *World Health Organization, Geneva*.

European Topic Centre on  
Human health and the environment  
<https://www.eionet.europa.eu/etcs/etc-he>

The European Topic Centre on Human health and  
the environment (ETC-HE) is a consortium of  
European institutes under contract of the European  
Environment Agency.

